

Snakes from the Pontal do Triângulo Mineiro Region, Cerrado Domain, Southeastern Brazil

Lucas Henrique Carvalho Siqueira¹ e Kátia Gomes Facure²

¹Instituto Butantan, Laboratório de Ecologia e Evolução, São Paulo, SP, Brasil.
hlucascs@gmail.com

²Faculdade de Ciências Integradas do Pontal, Universidade Federal de Uberlândia, Ituiutaba, MG, Brasil.
katiafacure@gmail.com

ABSTRACT:

Snakes play an important ecological role, interacting with other species and feeding on some that are considered pests and illness vectors. The present work aims to list the snake species present in the Municipality of Ituiutaba, southeastern Brazil. The data were obtained from preserved specimens deposited in zoological collections, visual surveys and occasional encounters. Fifty-seven individuals were analyzed, and we identified twenty-three species, distributed in five families. The number of species found is relatively high, considering the advanced degree of alterations in the natural environment, where only 15% remains intact. In our sample, we identified individuals of two viperid genera with medical and pharmacological importance. The present study contributes to the knowledge on snake assemblages in the Cerrado, and emphasizes the importance of inventories of the local fauna, mostly in environments with anthropic action, resulting in loss of habitats.

Keywords: *Species. Snakes. Ituiutaba.*

1 Introduction

Snakes are a highly diverse group, which are currently 392 known species in Brazil (COSTA and BÉRNILS, 2015). The Cerrado biome harbors a large number of these species (COSTA et al. 2007; MARQUES et al. 2015), even though herpetofauna is still poorly documented in several localities (COSTA et al. 2007). In five years the number of snake species rose from 107 in 2002 to 145 in 2007 (COLLI et al. 2002; COSTA et al. 2007), which reinforces the high diversity of snakes in this biome.

Agricultural expansion and intensification, together with cattle raising and urban development observed in the last decades in the Cerrado, the diversity of snakes may have been mainly affected by the loss and fragmentation of natural habitats (PONTES et al. 2009; WADE et al. 2003). This group has great ecological significance and the consequences of diversity loss can be severe, because as predators, they contribute to prey population control (ROCHA et al. 2008).

Some species in the Cerrado are considered as dangerous because of their highly toxic venom, which has medical and pharmacological importance, and the epidemiological distribution of accidents involving them is deeply linked with habitat modifications (WEN et al. 2002). Therefore, habitat loss may cause the snakes to invade anthropic environments, increasing the number of accidents (BASTOS et al. 2005) and frequently being killed by local residents.

The Cerrado biome is a recognized global biodiversity hotspot, with at least 87 priority areas for conservation (CAVALCANTI and JOLY, 2002). The Triângulo Mineiro region, in the west of the state of Minas Gerais, is especially under pressure due to the increase in cultivated areas, mainly for sugar cane, which is one of the major threats to biodiversity and can lead to population decline and local extinctions (GIARETTA et al. 2008; MICHELOTTO, 2008).

Species composition in a determined area is important as a framework for future research related to biodiversity and conservation (COLLI et al. 2002), thus, the aim of the present work was to inventory the snake species of the municipality of Ituiutaba, Minas Gerais. In addition, we compare the species list with a previous snake species list for the municipality of Uberlândia.

2 Materials And Methods

Study site

The municipality of Ituiutaba is located in the “Pontal do Triângulo Mineiro”, in the west of the State of Minas Gerais, (49° 27' 54" W; 18° 58' 08" S, figure 1), with a total area of about 2600 km². The local biome is characterized mainly by Cerrado phytophysionomies (IBGE, 2016). The climate is tropical with rainy summers and dry winters, where the average monthly temperature is about 18°C (TONIETTO, 2006), and an average annual rainfall of 1450 millimeters (QUEIROZ, 2012).



Figure 1 - Location of the Municipality of Ituiutaba (in yellow) in the state of Minas Gerais (in green), southeastern Brazil

Data collection

The work was carried out between May 2011 and June 2012 in non-standardized campaigns. Data collection was made through visual surveys for snakes in the field, on trails, access roads and forest patches near the university in Ituiutaba (Faculdade de Ciências Integradas do Pontal; FACIP - UFU), during diurnal and nocturnal periods. Possible shelters, such as fallen logs and burrows, were surveyed, as well as in vegetation up to five meters high (ZANELLA and CECHIN, 2006). Snakes found during the field activities of the FACIP students and teachers were identified and added to the list.

Snakes from the Zoonosis Control Center (CCZ) collection were identified and added to the list as well. Additional records were obtained by consulting the herpetological collections of the João Moojen Zoological Museum (MZUFV) at the Federal University of Viçosa, and the Ezequiel Dias Foundation (FUNED).

The encounters with snakes were photographically documented, however, they were not captured. Dead animals we encountered were deposited in the FACIP teaching collection. All snakes were identified to the lowest taxonomic level based on identification keys according to Peters and Orejas-Miranda (1970) and Campbell and Lamar (2004), comparing the specimens with photographic guides: Marques et al. (2004) and Marques et al. (2015). The resulting list was then compared to a previously published snake species list from the Municipality of Uberlândia, located about 130 kilometers from Ituiutaba (BRITES and BAUAB, 1988). The species names presented here follow those of Costa and Bérnils (2015).

3 Results

During the study, 57 specimens were examined. We recorded for the “Pontal do Triângulo Mineiro” region 23 snake species, distributed in 16 genera and five families (Table 1, Figure 2). In field surveys, (FS) we registered 17 species, including all species of the Boidae family with occurrence in the region. In the Zoonosis Control Center (CCZ) collection, 13 species were identified (six species were only sampled at CCZ), two species were recorded at Ezequiel Dias Foundation (FUNED), all previously recognized in our sample (visual surveys and occasional encounters), and one new record at João Moojen Zoological Museum (MZUFV): *Xenodon merremii*. The snakes were classified in four categories based on the state in which they were found: Preserved (n=33), Alive (n=9), Road-killed (n=13) or Killed by humans (n=2), (Figure 3).

Table 1: List of snake species recorded in the municipality of Ituiutaba, with the number of individuals (N), and origin: Field Survey (FS), Zoonosis Control Center (CCZ), Ezequiel Dias Foundation (FUNED), and João Moojen Zoological Museum of the Federal University of Viçosa (MZUFV). The (*) represents the species recorded in the municipality of Uberlândia.

Family/Species	FS	CCZ	FUNED	MZUFV
Boidae				
<i>Boa constrictor</i> Linnaeus, 1758*	1			
<i>Epicrates cenchria</i> (Cope, 1862)	1			
<i>Eunectes murinus</i> (Linnaeus, 1758)	2			
Colubridae				
<i>Chironius flavolineatus</i> (Boettger, 1885)*		2		
<i>Mastigodryas bifossatus</i> (Raddi, 1820)*		2		
<i>Simophis rhinostoma</i> (Schlegel, 1837)*	1			
Dipsadidae				
<i>Apostolepis assimilis</i> (Reinhardt, 1861)*		3	1	
<i>Erythrolamprus almadensis</i> (Wagler in Spix, 1824)*	1			
<i>Erythrolamprus poecilogyrus</i> (Wied, 1825)*	1			
<i>Erythrolamprus reginae</i> (Linnaeus, 1758)*	1			
<i>Oxyrhopus guibei</i> Hoge & Romano, 1978*	2	2	2	
<i>Oxyrhopus trigeminus</i> Duméril, Bibron & Duméril, 1854*		1		
<i>Philodryas agassizii</i> (Jan, 1863)*	1			
<i>Philodryas nattereri</i> Steindachner, 1870		1		
<i>Philodryas olfersii</i> (Lichtenstein, 1823)*	1	1		

to be continued...

Table 1: continuation...

Family/Species	FS	CCZ	FUNED	MZUFV
<i>Philodryas patagoniensis</i> (Girard, 1858)*		1		
<i>Phimophis guerini</i> (Duméril, Bibron & Duméril, 1854)*	1	1		
<i>Sibynomorphus mikanii</i> (Schlegel, 1837)*	3	5		
<i>Xenodon merremii</i> (Wagler in Spix, 1824)*				1
Elapidae				
<i>Micrurus frontalis</i> (Duméril, Bibron & Duméril, 1854)*	1			
Leptotyphlopidae				
<i>Trilepida koppesi</i> (Amaral, 1955)	2			
Viperidae				
<i>Bothrops moojeni</i> Hoge, 1966*	3	3	4	
<i>Bothrops neuwiedi</i> (Wagler, 1824)*	1	1		
<i>Crotalus durissus</i> (Linnaeus, 1758)*	1	1	1	
Total number of individuals	24	24	8	1
Total number of species	17	13	4	1



Figure 2: Representatives of snake species found during the field study: A) *Epicrates cenchria*; B) *Erythrolamprus poecilogirus*; C) *Erythrolamprus reginae*; D) *Phimophis guerini*; to be continued....



Figure 2: continuation...E) *Sibynomorphus mikanii*; F) *Bothrops moojeni*; G) *Bothrops neuwiedi*; H) *Crotalus durissus*. Photography authors: Siqueira, L. H. C. (B, C and E) and Facure, K. G. (A, D, F, G and H).

The Dipsadidae was the most diverse family, corresponding to 56.5% of the total richness and 55% of the total abundance. Considering all individuals analyzed, the most common species was *Bothrops moojeni* (n=10, 18.5%), followed by *Sibynomorphus mikanii* (n=8, 14.8%), and *Oxyrhopus guibei* (n=6, 11%). Four analyzed individuals could not be identified, three because of the poor state of conservation and roadkill in one case.

Twenty-eight species were recorded in the municipality of Uberlândia, being that nineteen species of that list were found in this study (Table 1). Nine species (*Liotyphlops ternetzii*, *Clelia clelia*, *Erythrolamprus aesculapi*, *Oxyrhopus rhombifer*, *Pseudoboa nigra*, *Sibynomorphus turgidus*, *Thamnodynastes strigilis*, *Micrurus lemniscatus*, and *Bothrops alternatus*) were not registered in our study. On the other hand, four species were recorded only in the municipality of Ituiutaba (*Epicrates cenchria*, *Eunectes murinus*, *Philodryas nattereri*, and *Trilepida koppesi*).




Figure 3 - Number of snakes found in each category: Preserved (Pr), Alive (Al), Road-killed (RK), and Killed by Humans (KH)

4 Discussion

The snakes we found have already been reported in the Cerrado and correspond to about 15.6% of the richness of this biome and none endemic species were recorded (COLLI et al. 2002; COSTA, 2007). The number of species found was smaller than those found in well-preserved and protected natural areas in this biome (richness ranged from 29 to 47; ARAUJO et al. 2010; FREITAS et al. 2016; RODRIGUES et al. 2015; SAWAYA et al. 2008; VALDUJO et al. 2009; VAZ-SILVA et al. 2007). However, considering the highly disturbed areas in the region of Ituiutaba, the snake assemblage presented here is relatively diverse. Besides, the visual surveys were conducted during the dry season (April, May and June), whereas snakes tend to have their activity peak during the wet/rainy season (BROWN et al. 2002; BROWN and SHYNE, 2002; MARQUES et al. 2000).

Five species (27.7%) from our sample are considered potentially dangerous (*Philodryas olfersii*, *Micrurus frontalis*, *Bothrops moojeni*, *Bothrops neuwiedi* and *Crotalus durissus*), including the most abundant species, *Bothrops moojeni*. This number represents almost one third of the species sampled in this region, which draws attention to the need of studies of these species mostly for the medical area (BERNARDE, 2011). Still, considering the proximity to the urban environment, several snakes can feed on animals that possibly are illness vectors, such as rodents, bats and snails (ZANOTTI-MAGALHÃES et al. 1991).

Several taxonomic groups have their historical classification marked by confusion, and the *Bothrops* complex is a very good example of it. During the data collection of the present study, the genus *Bothrops* was divided into “*Rhinocerophis*, *Bothrops* and *Bothropoides* (*Bothrops pauloensis* was included in this list)” (FENWICK et al. 2009). Carrasco et al. (2012) proposed the synonymization of the three genera under *Bothrops* a few years later. This instability in nomenclature is a complicating factor to taxonomy and future surveys and can be especially dangerous for medical and pharmacological issues (CARRASCO et al. 2016).

There were a huge number of snakes found dead either roadkill or killed by humans (Figure 3). This number must be due to the length of the snakes and their slow speed when crossing roads, besides they have the habit of using these sites, which provides considerable heat for thermoregulation (KUNZ and GHIZONI Jr., 2009).

Comparing our results with those shown by Brites and Bauab (1988) for the municipality of Uberlândia, it is possible to observe that the number of sampled species is approximately the same. This fact can be explained by the proximity of the study sites, despite the divergence in time expended in data collection (13 months in this work and 39 months in BRITES and BAUAB, 1988) and methodology (field surveys and material examined in collections in this work, while snakes were brought to the university by citizens in BRITES and BAUAB, 1988). Combining both lists, the total number of snake species known to the Triângulo Mineiro region is thirty-two. In both studies, *S. mikanii* was one of the most abundant, indicating that this species is very common in that region.

The species richness in the municipality of Ituiutaba is probably higher, and this subsample might be associated partly with the fossorial or cryptic habits of some snakes, such as *Apostolepis assimilis* (LEMA, 2001), which makes it difficult to find those species. Such habits can explain the absence of snakes of the Anomalepididae family, recorded previously in Uberlândia (BRITES and BAUAB, 1988).

5 Conclusions

The snake assemblage of the municipality of Ituiutaba has a large diversity of habitat use, and many are widely distributed, including other biomes, e.g. the Atlantic Forest (MARQUES et al. 2004). Hence, it is necessary to consider that due to strong anthropic pressure, since there are different ways the species may be impacted, mainly by population decline and community homogenization (MCKINNEY, 2006). It is important to stress that the majority of snakes recorded were identified from preserved animals in collections; such fact is an evidence of the great importance of zoological collections for biodiversity studies.

For the species discussed here, many have ecological and medical importance (ROCHA et al. 2008), which evidences the necessity to know the local fauna. The present study contributes to the knowledge about the snake assemblage in the west of Minas Gerais (Ituiutaba region), and sheds light on the need of future studies on diversity and distribution of this vertebrate group in the Cerrado under an unfavorable scenario with respect to conservation of natural areas.

Acknowledgements

We are grateful to the Zoonosis Control Center of Ituiutaba for specimen loans for this study, and to MZUFV and FUNED for the access to specimens. We thank all the people who contributed to the data collection, especially Ariovaldo A. Giaretta and Thaís de Cássia Rodrigues. To Paulo Sérgio Bernarde for helping with species identification, to all friends and staff from FACIP - UFU that collaborated on this work. We thank Thiago R. de Carvalho for critically reading an earlier version and providing valuable insight into this manuscript, and for linguistic help.

Referências

- ARAUJO CO, CORRÊA-FILHO DT, SAWAYA RJ. Snake Assemblage of Estação Ecológica de Santa Bárbara, SP: a Cerrado Remnant in Southeastern Brazil. *Biota Neotrop.* 2010;10(2):235-245.
- BASTOS EGM, ARAUJO AFB, SILVA HR. Records of the rattlesnakes *Crotalus durissus terrificus* (Laurenti) (Serpentes, Viperidae) in the State of Rio de Janeiro, Brazil: a possible case of invasion facilitated by deforestation. *Rev. Bras. Zool.* 2005;22(3):812-815.
- BERNARDE PS. Mudanças na classificação de serpentes peçonhentas brasileiras e suas implicações na literatura médica. *Gaz. Méd. Bahia.* 2011;81(1):55-63.
- BROWN GP, SHINE R. Influence of weather conditions on activity of tropical snakes. *Austral Ecol.* 2002;27:596-605.
- BROWN GP, SHINE R, MADSEN T. Responses of three sympatric snake species to tropical seasonality in Northern Australia. *J. Trop. Ecol.* 2002;18:549-568.
- BROWN GP, SHINE R, MADSEN T. Spatial ecology of slatey-grey snake (*Stegonotus cucullatus*, Colubridae) on a tropical Australian floodplain. *J. Trop. Ecol.* 2005;21:605-612.
- BRITES VL, BAUAB FA. Fauna Ofidiana do Município de Uberlândia, Minas Gerais – Brasil I. Ocorrência na Área Urbana. *R. Cent. Ci. Bioméd. Univ. Fed. Uberlândia.* 1998;4(1):3-8.
- CAMPBELL JA, LAMMAR WW. The venomous reptiles of the Western Hemisphere, 1st ed. New York: Ithaca; 2004.
- CARRASCO PA, MATTONI CI, LEYNAUD GC, SCROCCHI GJ. Morphology, phylogeny and taxonomy of South American Bothropoid pitvipers (Serpentes: Viperidae). *Zool. Scr.* 2012;41(2):109-124.
- CARRASCO PA, VENEGAS PJ, CHAPARRO JC, SCROCCHI GJ. Nomenclatural instability in the venomous snakes of the *Bothrops* complex: implications in toxinology and public health. *Toxicon.* 2016;119:122-128.
- CAVALCANTI RB, JOLY CA. Biodiversity and conservation priorities on the Cerrado Region. In: OLIVEIRA P, MARQUIS RJ, editors. *The Cerrados of Brazil: ecology and natural history of a Neotropical Savanna.* New York: Columbia University Press; 2002. p. 351-367.
- COLLI GR, BASTOS RP, ARAUJO AFB. The character and dynamics of the Cerrado herpetofauna. In: OLIVEIRA P, MARQUIS RJ, editors. *The Cerrados of Brazil: ecology and natural history of a Neotropical Savanna.* New York: Columbia University Press; 2002. p. 223-241.
- COSTA HC, BÉRNILS RS. Répteis brasileiros: Lista de espécies 2015. *S. Am. J. Herpetol.* 2015;4(3):75-93.
- COSTA GC, NOGUEIRA CC, MACHADO RB, COLLI GR. Squamate Richness in the Brazilian Cerrado and its environmental-climatic associations. *Divers. Distrib.* 2007;13:714-724.
- FENWICK AM, GUTBERLET Jr. RL, EVANS JA, PARKINSON CL. Morphological and molecular evidence for phylogeny and classification of South American pitvipers, genera *Bothrops*, *Bothriopsis* and *Bothrocophias* (Serpente: Viperidae). *Zool. J. Linn. Soc-Lond.* 2009;156:617-640.
- FREITAS MA, COLLI GR, ENTIAUSPE-NETO OM, TRINCHÃO L, ARAÚJO D, LIMA TO et al. Snakes of Cerrado Localities in Western Bahia, Brazil. *Check List.* 2016;12(3):1-10.
- GIARETTA AA, MENIN M, FACURE GK, KOKUBUM MNC, FILHO JCO. Species richness, relative abundance, and habitat of reproduction of terrestrial frogs in the Triângulo Mineiro region, Cerrado biome, Southeastern Brazil. *Iheringia Ser. Zool.* 2008;98(2):181-188.

- IBGE - INSTITUTO BRASILEIRO DE GEOGRAFIA E ESTATÍSTICA [Internet] – Brasília: Ministério do Planejamento, Orçamento e Gestão (BR) [visited Mar 14, 2017]. Available from: <http://cidades.ibge.gov.br/xtras/perfil.php?codmun=313420>.
- KUNZ TS, GHIZONI Jr IR. Serpentes Encontradas Mortas em Rodovias do Estado de Santa Catarina, Brasil. *Biotemas*. 2009;22(2):91-103.
- LEMA T. Fossorial Snake Genus *Apostolepis* From South America (Serpentes: Colubridae: Elapomorphinae). *Cuad Herpetol*. 2001;15(1):29-43.
- MARQUES OAV, ETEROVIC A, ENDO W. Seasonal activity of snakes in the Atlantic Forest in Southeastern, Brazil. *Amphibia-Reptilia*. 2000;22:103-111.
- MARQUES OAV, ETEROVIC A, SAZIMA I. Snakes of The Brazilian Atlantic Forest, An Illustrated field guide for the Serra do Mar Range. 1 st ed. Ribeirão Preto: Holos; 2004.
- MARQUES OAV, ETEROVIC A, NOGUEIRA CC, SAZIMA I. Serpentes do Cerrado, guia ilustrado, 1 st ed. Ribeirão Preto: Holos; 2015.
- MICHELOTTO DB. Novos Arranjos Territoriais. A Expansão da Cultura da Cana-de-Açúcar na Região do Triângulo Mineiro [dissertation]. Brasília: Universidade de Brasília/UNB; 2008. 188p.
- MCKINNEY ML. Urbanization as a major cause of biotic homogenization. *Biol. Conserv*. 2006;127:247-260.
- PETERS JA, OREJAS-MIRANDA B. Catalogue of the Neotropical Squamata: part I. snakes. *U. S. Nat. Museum Bull*. 1970;297:297-347.
- PONTES JAL, PONTES RC, ROCHA CFD. The snake community of Serra do Mendanha in Rio de Janeiro state, Southeastern Brazil: composition, abundance, richness and diversity in areas with different conservation degrees. *Braz. J. Biol*. 2009;69(3):795-804.
- QUEIROZ AT, COSTA RA. Caracterização e variabilidade climática em séries de temperatura, umidade relativa do ar e precipitação em Ituiutaba-MG. *Caminhos da Geografia*. 2012;13(43):346-357.
- ROCHA CFD, BERGALLO HG, VERA Y CONDE CF, BITTENCOURT EB, SANTOS HC. Richness, abundance, and mass in snake assemblages from two Atlantic Rainforest sites (Ilha do Cardoso, São Paulo) with differences in environmental productivity. *Biota Neotrop*. 2008;8(3):117-122.
- RODRIGUES JB, GAMA SCA, FILHO GAP, FRANÇA FGR. Composition and ecological aspects of a snake assemblage on the Savanna Enclave of the Atlantic Forest of the Guarbas Biological reserve in Northeastern Brazil. *S. Am. J. Herpetol*. 2015;10(3):157-164.
- SAWAYA RJ, MARQUES OAV, MARTINS M. Composition and natural history of a Cerrado snake assemblage at Itirapina, São Paulo, Southeastern Brazil. *Biota Neotrop*. 2008;8(2):127-149.
- TONIETTO J, VIANELLO RL, REGINA MA. Caracterização macroclimática e potencial enológico de diferentes regiões com vocação vitícola de Minas Gerais. *Informe Agropecuário, Belo Horizonte*. 2006;27(234):32-55.
- VALDUJO PH, NOGUEIRA CC, BAUMGARTEN L, RODRIGUES FHG, BRANDÃO RA, ETEROVIC A, et al. Squamate reptiles from Parque Nacional das Emas and surroundings, Cerrado of Central Brazil. *Check List*. 2009;5(3):405-417.
- VAZ-SILVA W, GUEDES AG, AZEVEDO-SILVA PL, GONTIJO FF, BARBOSA RS, ALOÍSIO GR et al. Herpetofauna, Espora Hydroelectric Power Plant, state of Goiás, Brazil. *Check List*. 2007;3(4):338-345.
- WADE TG, RIITTERS KH, WICKHAN JD, JONES KB. Distribution and causes of global forest fragmentation. *Conserv. Ecol*. [Internet]. 2003 [cited 2017 mar 14];7(2):7. Available from: <http://www.consecol.org/vol7/iss2/art7/>

WEN FH, CARDOSO JLC, MÁLAQUE CMS, FRANÇA FOS, SANT'ANNA SS, FERNANDES W, et al. Influência das Alterações Ambientais na Epidemiologia dos Acidentes Ofídicos e na Distribuição Geográfica das Serpentes de Importância Médica nos Estados de São Paulo e Paraná, 1988-1997. Informe Epidemiológico do SUS. 2002;11(1):45-47.

ZANELLA N, CECHIN SZ. Taxocenose de Serpentes no Planalto Médio do Rio grande do Sul, Brasil. Rev. Bras. Zool. 2006;23(1):211-217.

ZANOTTI-MAGALHÃES EM, MAGALHÃES LA, DE CARVALHO JF. Relação entre patogenicidade de *Schistosoma mansoni* em camundongos e susceptibilidade do molusco vetor. I. Infeciosidade das Cercárias e cargas de vermes. Rev. Saude Publ. 1991;25(5):359-366.