

## Physicochemical properties, toxicity and larvicidal activity of the essential oil of *Cymbopogon winterianus* in front of *Aedes Aegypti*

Gabriela Morais da Costa<sup>I</sup>, Maria Conceição Carneiro dos Santos<sup>II</sup>, Gustavo Oliveira Everton<sup>III</sup>, Paulina de Cássia Duarte de Sousa<sup>IV</sup>, Maurício Eduardo Salgado Rangel Mail<sup>V</sup>

### ABSTRACT

Due to the increasing rate of morbidity and mortality caused by vector diseases in the current context, especially by *Aedes aegypti*, substances of plants have been studied as alternatives to chemical insecticides, among them, the essential oil (EO) of the species *Cymbopogon winterianus*. Thus, this study evaluated the larvicidal activity of the EO of *C. winterianus* against the *A. aegypti* larvae. EO was extracted through the hydro-distillation technique and physicochemical properties were determined. To evaluate larvicidal activity, tests were performed with larvae in the third instar at the final concentrations of 19.54; 26.50; 55.59; 138.98; 208.47 and 277.97 mg L<sup>-1</sup> of *C. winterianus* EO. In addition, *Artemia salina* Leach bioassay was used to verify toxicity effect. EO obtained presented satisfactory results in 2.64%. In the larvicidal assay, 100% mortality of larvae was observed after 24 hours at concentrations of 208.47 mg L<sup>-1</sup> and 277.97 mg L<sup>-1</sup> of the EO, showed effective in the other concentrations and with LC50 of 46.18 mg L<sup>-1</sup>, considered highly active. In the toxicity assay, the EO presented LC50 at 532.34 mg L<sup>-1</sup>, showed considered nontoxic. These results reinforce the use of EO front *A. aegypti* larvae control.

**Keywords:** Arboviruses control; Hydro-distillation; Biological assays

### RESUMO

Devido a crescente taxa de morbimortalidade causada por doenças vetoriais no contexto atual, sobretudo pelo *Aedes aegypti*, substâncias de plantas vêm sendo estudadas como alternativas aos inseticidas químicos, entre elas, o óleo essencial (OE) da espécie *Cymbopogon winterianus*. Desta forma, este estudo avaliou a atividade larvicida do OE de *C. winterianus* frente as larvas do mosquito *Aedes aegypti*. O OE foi extraído através da técnica de hidrodestilação e parâmetros físico-químicos foram determinados. Para avaliação da atividade larvicida foram realizados testes com larvas no terceiro estágio nas concentrações finais de 19.54; 26.50; 55.59; 138.98; 208.47 e 277.97 mg L<sup>-1</sup> do óleo essencial de *C. winterianus*. Além disso, o bioensaio de *Artemia salina* Leach foi utilizado para verificação da toxicidade. O OE obtido apresentou rendimento satisfatório em 2,64%. Nos ensaios larvicidas observou-se mortalidade de 100% das larvas após 24 horas nas concentrações 208,47 mg L<sup>-1</sup> e 277,97 mg L<sup>-1</sup> do OE, sendo eficazes nas demais concentrações e com CL50 de 46,18 mg L<sup>-1</sup>, considerado altamente ativo. No ensaio de toxicidade, o OE apresentou CL50 em 532,34 mg L<sup>-1</sup>, sendo considerado atóxico. Estes resultados reforçam o uso do óleo essencial frente ao controle de larvas de *A. aegypti*.

**Palavras-chaves:** Controle de arbovíroses; Hidro-destilação; Ensaios biológicos

<sup>I</sup> Universidade Federal do Maranhão, São Luís, MA, Brasil - gabi\_costa\_10@hotmail.com

<sup>II</sup> Universidade Federal do Maranhão, São Luís, MA, Brasil - mariaccscontato@hotmail.com

<sup>III</sup> Universidade Federal do Maranhão, São Luís, MA, Brasil - gustavooliveiraeverton@gmail.com

<sup>IV</sup> Universidade Federal do Maranhão, São Luís, MA, Brasil - paulinaduarte000@gmail.com

<sup>V</sup> Universidade Federal do Maranhão, São Luís, MA, Brasil - mauricio.rangel@ufma.br



## 1 INTRODUCTION

*Aedes aegypti* is a vector of the most widespread urban arboviruses worldwide, causing high morbidity and mortality such as dengue, chikungunya, zika and yellow fever, especially in countries where environmental, climatic and sanitary conditions favor its reproduction (LIMA-CAMARA, 2016). Transmission of vector diseases occurs when the infected female of *A. aegypti* encounters, during daytime and low flights, a susceptible host and transmits the arbovirus through saliva while sucking up the blood, the latter it is essential for egg maturation. After this, the pregnant female seeks reservoirs with clean water and stops to begin oviposition; laying between 150 to 200 eggs which will start the larval phase of the mosquito upon hatching, consisting of 4 instars over a duration of 10 days. Upon entering adulthood, adult mosquitos mate and to trigger a new cycle period with an interval of 30 to 35 days of adulthood (CANTANE et al., 2015).

The Ministry of Health from Brazil adopts different measures to prevent vector-borne diseases. The practice of chemical control mechanisms is widely used to combat the mosquito *A. aegypti*. It occurs mainly through the use of synthetic insecticides that release neurotoxins, analogous juvenile hormones and chitin synthesis inhibitors capable of killing mosquitoes in their larval and adult stages (ZARA et al., 2016). However, the frequent and intensive use of these chemicals causes the emergence of resistant *A. aegypti* populations, in addition to the undesirable effects on the ecosystem, and supporting species for environmental balance due to the lack of selectivity and long-term effects of those products on the environment (CORRÊA, 2011; VELOSO et al., 2015).

The application of plants compounds as an alternative method to the use of insecticides has numerous advantages such as rapid degradation, low resistance potential, high availability and high economic viability (CORRÊA, 2011). Essential oil from plants is obtained by releasing volatile compounds encapsulated in "pouches" called trichomes, which are produced for self-defense and attraction of pollinators and are found in various parts of the plant, as leaves, flowers and stalks

(WOLFFENBUTTEL, 2007). Natural compounds have been investigated by several researchers who had proven the potential larvicide of plants against different species of insects, including *A. aegypti* (CARVALHO et al., 2003; CAVALCANTI et al., 2004; CHENG et al., 2004; MORAIS et al., 2006; PUSHPANATHAN et al., 2006).

*Cymbopogon* spp, popularly known as citronella grass, is an aromatic and perennial plant with long, flat leaves and curved appearance, which is grown in tropical regions for use in preparation of teas and for its known antifungal, antibacterial and insect repellent properties that are obtained by extracting its essential oils (VELOSO et al., 2015). Essential oil (EO) from citronella grass (*C. winterianus*) demonstrates good performance as a natural repellent and larvicide, as its chemical constitution is integrated with major components such as aldehyde, citronellal, geraniol and citronellol, to which therapeutic, pharmaceutical and cosmetic efficiency characteristics are attributed (CASSEL et al., 2009; KAKARAPARTHI et al., 2014). Studies developed on the efficacy of the use of *C. winterianus* essential oil has been recommended as an alternative in arboviruses control, as a way to improve its possibility of use (MACEDO et al., 2010). Thus, this study evaluating the toxicity and larvicidal activity of the EO of *C. winterianus* (citronella grass) against the larvae of *Aedes aegypti*.

## 2 METHODOLOGY

*C. winterianus* leaves were harvested on 9th January, 2020 between 10 a.m. and 12 p.m. at the Berta Lange Morretes Medicinal Plants Garden, located at Federal University of Maranhão (UFMA), São Luís, Maranhão, Brazil and identified by specialists Prof. Dr. Ana Zélia Silva. The leaves was rinsed, ground (cut into small parts) and dried in greenhouse at 45° C for 24 h.

Leaves was subjected to the hydro-distillation technique using a glass Clevenger extractor coupled with a round bottom flask packed in an electric blanket as a heat source. 300 g of *C. winterianus* leaves were used, adding distilled water (1:10). Hydro-distillation was conducted at 100°C for 3 hours and the extracted EO was collected.

The oil was dried over anhydrous sodium sulfate ( $\text{Na}_2\text{SO}_4$ ), and was then stored in glass bottles covered with aluminum foil and stored in a refrigerator ( $4^\circ\text{C}$ ) until further investigation. The yield of essential oil was expressed in percentage w/w.

Physicochemical properties such as density, refractive index, solubility, color and odor were determined according to the Brazilian Pharmacopoeia (2019).

A. aegypti trap and ovitrap, adapted from FAPERJ (2013), were used to capture the larvae. These systems consisted of a 1 gallon PET bottle, cut in half allowing to reach a larger opening area, where 1 mL of an aqueous solution containing a concentration of 12% (6 g / 50 mL) brewer's yeast was aspirated and added to 300 mL of running water and black polyethylene pails (50 mL) filled with chlorinated water and two eucatex vanes to oviposition.

The traps installed were sheltered from the sun and rain, in neighborhoods with a high occurrence of the vector in the capital São Luís of State of Maranhão, were inspected weekly and analyzed in the Laboratory of Research and Application of Essential Oils (LOEPAV/UFMA).

The larvae collected were kept in plastic trays containing clean water with pH 6.9, and remained for a photoperiod of 14 hours of light and 10 hours of darkness in laboratory conditions ( $29 \pm 1^\circ\text{C}$ ), being fed with finely ground cat food until they reached the 3rd instar (L3).

The biological assays were performed according to the methodology recommended by the WHO (1970) adapted by Veloso et al. (2015). The larvae that reached the 3rd instar were separated by means of a pipette, placed on filter paper for the removal of excess water and then 10 larvae were deposited in beakers containing 30 mL of distilled water, where EO aliquots diluted in 500  $\mu\text{L}$  of dimethyl sulfoxide (DMSO) were added to formalize the final concentrations of 19.54; 26.50; 55.59; 138.98; 208.47 and 277.97  $\text{mg L}^{-1}$  corresponding to the aliquots of 0.70 $\mu\text{L}$ /500 DMSO, 0.95  $\mu\text{L}$ /500 DMSO, 2.0  $\mu\text{L}$ /500 DMSO, 5.0  $\mu\text{L}$ /500 DMSO, 7.5  $\mu\text{L}$ /500 DMSO e 10  $\mu\text{L}$ /500 DMSO of *C. winterianus* essential oil, respectively. The bioassays were performed in triplicate.

Mortality was verified after 24 h of exposure concentrations in EO. The larvae's death was declared by absence of muscle contraction when stimulated with a pipette.

Considering the percentage of mortality for each concentration, the median lethal concentration (LC50) was submitted to Probit analysis (log concentrations), with a 95% confidence interval, using the software R version 3.6.3 or Excel version 2010.

The toxicity test was carried out according to the methodology described by Meyer *et al.* (1982). In a rectangular container, with a partition containing holes of approximately 0.02 cm in diameter, spaced by 0.5 cm and evenly distributed, artificial saline solutions (60 g L<sup>-1</sup> of distilled water) (60 g of sea salt/liter of distilled water) were added. The container was placed inside an incubator illuminated by a fluorescent lamp, with aeration. On one side of this container, about 64 mg of *A. salina* cysts were added, given that they did not cross the partition. The part of the system containing *A. salina* cysts was covered with aluminum foil, so that the organisms, at birth, were attracted by light on the other side of the system, forcing them to cross the partition. This procedure aims to homogenize the physical conditions of the test organisms. Incubation was performed for a period of 48 hours. The temperature was monitored during all bioassay.

For the evaluation of the lethality of *A. salina*, a stock saline solution of EO was prepared at a concentration of 10,000 mg L<sup>-1</sup> and 0.02 mg of Tween 80 (surfactant). From this, aliquots of 5, 50 and 500 µL were transferred to test tubes and completed by adding saline solution previously prepared up to 5 mL, obtaining concentrations of 10, 100 and 1000 mg L<sup>-1</sup>, respectively. All tests were carried out in triplicates, where 10 larvae in the nauplii phase were transferred to each of the test tubes.

For control 5 mL of saline solution was used, for positive control K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> and for negative control 5 mL of a solution 4 mg L<sup>-1</sup> of Tween 80. After 24 hours of exposure, the live larvae were counted by considering exclusively those that did not move during observation period or only with the slight agitation of the vial. In order to classify the EO toxicity, the criterion established by Dolabela (1997) was adopted; being considered a highly toxic product when LC50 ≤ 80 mg L<sup>-1</sup>, moderately

toxic to  $80 \text{ mg L}^{-1} \leq \text{LC50} \leq 250 \text{ mg L}^{-1}$  and mildly toxic or nontoxic when  $\text{LC50} \geq 250 \text{ mg L}^{-1}$ .

### 3 RESULTS AND DISCUSSION

The physicochemical properties obtained for the *C. winterianus* EO are presented in Table 1.

Table 1 – Physicochemical properties of the essential oil of *C. winterianus* leaves.

Physicochemicals properties	<i>C. winterianus</i> EO
Density ( $\text{g mL}^{-1}$ )	0,85
Solubility in ethanol 70 % (v/v)	1:6
Solubility in ethanol 90 % (v/v)	1:1
Yield (w/w %)	2,64
Refractive index at 25° C	1,475
Color	Yellow
Odor	Characteristic

The density of the extracted EO is less than water ( $1 \text{ g mL}^{-1}$ ). The extracted EO presented a yellowish color, with a clear appearance and strong odor. It was found that the EO in question has better solubility in interaction with 90% ethanol, based on proportionate comparison to a concentration of 70%, although both are considered easily soluble according to the Brazilian Pharmacopoeia. The percentage yield obtained through hydro-distillation was 2.64%, expressing a mass of 2.64 g EO extracted for every 100 g of dried leaves of the species (Table 1).

According to the study by Pereira et al. (2016), in which the result in *Cymbopogon flexuosus* leaves collected in January was 3.73%, the EO yield is subject to monthly influence of the region temperature and soil moisture. According to Rocha et al. (2000), changes are due to variations in drying temperature, which, in comparison to their research, resulted in 0.987% drying at 40°C of leaves of *C. winterianus*.

The EO of the leaves of *C. winterianus* obtained in the same place of collection of the present study was investigated by Rodrigues et al. (2013), and presented yield

of 1.3%, without coloration and distinct odor: This distinction between the physicochemical properties analyzed may have been influenced by the harvest time. Based on the work developed by Blank et al. (2007), the yield of volatile citronella oils produced at 9:00 am by season with a value corresponding to 2.71, 2.36 and 4.24% for summer, winter and spring, respectively, as the volatile oil content generally peaked at that time.

The results of the larvicidal test for the L3 *Aedes aegypti* larvae in 6 distinct concentrations of 19.54; 26.50; 55.59; 138.98; 208.47 and 277.97 mg L<sup>-1</sup> of the *C. winterianus* EO are shown in Table 2.

Table 2 – Average mortality of *Aedes aegypti* larvae against the action of *Cymbopogon winterianus* essential oil

EO mg L <sup>-1</sup>	Nº dead	Nº alive	Deaths accumulated	Alives accumulated	Mortality (%)
277.97	10	0	39	0	100
208.47	10	0	29	0	100
138.98	9	1	19	1	90
55.59	5	5	10	6	50
26.50	3	7	5	13	30
19.54	2	8	2	21	20

Nº = average number of larvae per replicate

The EO was shown to be efficient in the control of larvae in all concentrations tested, being more promising in the concentrations of 208.47 mg L<sup>-1</sup> and 277.97 mg L<sup>-1</sup> where after 24 hours 100% of the larvae had died (Table 2). In the other concentrations, the mortality rate is below 90% in the 24-hour reading. Larvae in water and 2% DMSO showed null mortality during the test, indicating that the DMSO used for dilution of EO did not influence the larval death and biological activity in the treated groups. Considering that all EO concentrations of *C. winterianus* produced mortality data after 24 hours of exposure, it is possible to state that the plants show a high larvicidal effect in the larval stage of *Aedes aegypti*.

According to studies carried out by Amer and Mehlhorn (2006), the EO of *C. winterianus* caused 60% mortality in *A. aegypti* larvae after 24 hours of exposure to a concentration of 50 mg L<sup>-1</sup> similar to the action of EO at the concentration of 55.59 mg

L<sup>-1</sup> presented in this study. On the other hand, the larvicidal assays developed by Falcão (2018) also reached the maximum mortality percentage (100%) of *A. aegypti* with an esterified (250 mg L<sup>-1</sup>) and crude (250 mg L<sup>-1</sup>) EO of *C. winterianus*, however, are higher concentration values than those observed in this study. This fact may have occurred due to several factors such as origin, seasonality, extraction method, collection and synergism of the major chemical compounds of the species as citronellal, citronelol and geraniol that are the main responsible for larvicidal activity (GOMES et al., 2019; LEITE et al., 2011; SINHA et al., 2014).

In the study developed by Veloso et al. (2015), the larvicidal effect of citronella grass EO caused 100% of the mortality of *A. aegypti* larvae from the aliquot of 10 µl, while the aliquots of 5,0 µl and 7,5 µl presented the same result after 6 hours of exposure from the beginning of the experiment.

*Cymbopogon nardus* belonging to the same genus of plant evaluated in this work and its EO was investigated by Phasomkusolsil and Soonwera (2010). The authors found that EO presented 100% mortality in 3rd instar larvae of *Culex quinquefasciatus* and *Anopheles minimus* species after 5 and 10 minutes of exposure, respectively. The results of this study corroborate the efficacy of plant species of the genus *Cymbopogon* against vector insects, proving to be a promising strategy for moderating the indiscriminate use of synthetic insecticides that has promoted the emergence of resistant populations (Veloso et al., 2015).

Statistical analysis indicated that the lethal concentration capable of reducing 50% of the larvae with the EO was 46.18 (41.62 - 50.74) mg L<sup>-1</sup> (Table 3).



Table 3 - Larvicidal activity of *Cymbopogon winterianus* essential oil against cultivated larvae of *Aedes aegypti*.

Exposure time	Concentration mg L <sup>-1</sup>	Mortality (%)	LC50 <sup>a</sup> (mg L <sup>-1</sup> ) (IC 95%)
24 hours	DMSO	0	46,18 (41,62 – 50,74) <sup>b</sup>
	19,57	20	
	26,50	30	
	55,59	50	
	138,98	90	
	208,47	100	
	277,97	100	

a = The LC<sub>50</sub> was calculated by Probit analysis using the software R version 3.6.3

b = 95% confidence interval; no larvae killed in the negative control, composed of 2% DMSO solution; the positive control, 1 mg L<sup>-1</sup>

<sup>1</sup> temefos, exhibited 100% larval mortality

According to Cheng et al. (2003), the values of LC<sub>50</sub> < 50 mg L<sup>-1</sup> are considered highly active. Thus, from this classification, the EO extracted from the leaves of *C. winterianus* is considered of highly efficient potential, encouraging its application capacity.

The observed potential is justified by the *C. winterianus* being known to have insect repellent properties, being widely commercialized for this purpose (MENDONÇA et al., 2005). It is possible that the larvicidal potential of the EO against *A. aegypti* can be attributed to the presence of terpenic, alcohol and aldehyde components (LEE, 2006; GOMES et al., 2016).

Among the chemical larvicides used by the Ministry of Health from Brazil, pyriproxyfen is a field-applied development inhibitor for the treatment of domestic *A. aegypti* breeding sites and water for human consumption. In a bioassay recently performed by Fiaz et al. (2019), a LC<sub>50</sub> was estimated for pyriproxyfen equal to 8.20 mg L<sup>-1</sup>, where there were 6 concentrations of pyriproxyfen with triplicates containing 20 larvae (L3) of *A. aegypti* in 25 mL of distilled water. In addition to pyriproxyfen, when compared to other commercial products such as BTI (0.026 mg L<sup>-1</sup>) (FANSIRI et al., 2006), the efficacy of the EO obtained is considerably low, limiting its application in practice.

In comparison with the previous investigations of Mendonça et al. (2005), where the ethanol extract of *C. winterianus* leaves has a LC<sub>50</sub> value equal to 38.8 mg L<sup>-1</sup>

against *A. aegypti* larvae, the EO proved to be less efficient than the reported extract. However, for the others EOs that proposed the same bioassay in the literature, the EO exhibited higher larvicide action against *A. aegypti* larvae, such as EOs of *C. argyrophyllus* (LC50 = 310 mg L<sup>-1</sup>) (Cruz et al., 2017), leaves of *Mentha spicata* (LC50 = 56.08 mg L<sup>-1</sup>) (Govindarajan et al., 2011), roots of *Acorus calamus* L (LC50 = 99.41 mg L<sup>-1</sup>) (Manzoor et al., 2013), aerial parts of *Anethum graveolens* L. (LC50 = 50 mg L<sup>-1</sup>) (Amer and Mehlhorn, 2006), seeds of *Carum carvi* L (LC50 = 54.62 mg L<sup>-1</sup>) (Pitasawat et al., 2007), flowers of *Dendropanax morbifera* Leveille (LC50 = 62.32 mg L<sup>-1</sup>) (Chung et al., 2009) and fruits of *Heracleum pastinacifolium* subsp. *Transcaucasicum* (LC50 = 69.72 mg L<sup>-1</sup>) (Tabanca et al., 2012).

In previous studies, other biological activities of EO of *C. winterianus* were found, such as its acaricidal action investigated in larvae and adult females engorged with tick *Boophilus microplus* (Agnolin et al., 2014) and also molluscicide against adult snails of the genus *Biomphalaria*, being reported for the first time by studies conducted by Rodrigues et al., 2013. The insect repellent action has already been reported by several authors (MENDONÇA et al., 2005; EDEN et al., 2018), among these investigations the repellency of citronella oil associated with 5% vanilla to three species of mosquitoes, *Aedes aegypti*, *Culex quinquefasciatus* and *Anopheles* sp. who were exposed to EO for more than 8 hours (TAWATSIN et al., 2001). Regarding fungicidal activity, a study developed by Cruz et al., (2015) indicated a reduction in the mycelial growth of *F. solani* according to the addition of the oil under analysis.

In the United States of America or USA, the Federal Insecticide, Fungicide, and Rodenticide (FIFRA) is the federal law that establishes the basic system of pesticide regulation in the United States for registration, distribution, sale and use (EPA, 2020). According to ISMAN (2016), Section 25(b) of FIFRA, which contains chemicals with active and inert ingredients considered to be of minimal risk, includes a dozen essential oils, among them citronella oil (*C. winterianus*). In the European Union, citronella oil is being considered for approval for use as insect repellents.


Table 4 shows that the EO solution extracted from *C. winterianus* causes a 50% mortality rate in the *Artemia salina* Leach in concentrations  $1000 \text{ mg L}^{-1}$ , represented by the Log of dose 3.00.

Table 4 - Mortality of the species of *Artemia salina* Leach due to the action of essential oil

Log C	N° dead	N° alive	Deaths accumulated	Alives accumulated	Mortality (%)
3	5	5	5	8	50
2	3	7	12	3	30
1	0	10	22	0	0

The LC50 calculated by the intersection of the accumulated curve alive and dead presented in Figure 1 resulted in LC50 of  $532.34 \text{ mg L}^{-1}$  and by the Dolabela criterion (1997) this classifies the EO as non-toxic. This result is contrasting with that obtained by Brasileiro et al. (2006), who identified the EO of *Cymbopogon nardus* as toxic (LC50  $118.92 \text{ mg L}^{-1}$ ) for brine shrimp larvae. In addition, Pertiwi et al. (2014) investigated citronellal, one of the main chemical constituents of *C. winterianus*, performing its toxicity test against *A. salina* before and after 2 and 4 weeks of storage and found initial LC50 equal to  $71,1 \text{ mg L}^{-1}$ , showing to be highly toxic.

Figure 1- Curve of living and dead accumulations per concentration log



According to Ferreira et al. (2017), plants classified as toxic produce secondary metabolites that by inhalation, ingestion or contact can cause pathological changes in men and animals and, in certain situations, can lead to serious disorders in the body and even death.

Veletovac (2016) mentioned the importance of knowing the composition and concentration of the constituents of an EO, since its toxic potential is directly attributed. The author also reports that EOs of the genus *Cymbopogon* have a high range of application and have useful and popular substances, and after screening some EO of the family Poaceae, oils with large amounts of geraniol compared to those with citral as the main component can be classified as safe.

Thus, the obtained result is considered important, since the non-toxicity of the EO of *C. winterianus* proves its safety against non-target organisms in the regions of occurrence of insect vectors. No analyses were found to corroborate the non-toxicity of this oil in scientific productions, presenting an interesting characteristic for the use of EO studied in urban environments, where there is a higher incidence of water reservoirs that favor the proliferation of *A. aegypti*.

#### **4 CONCLUSION**

The essential oil of *Cymbopogon winterianus* presented larvicidal action against *Aedes aegypti*, proving to be more efficient for the aliquots of 7.5  $\mu\text{l}$  and 10.0  $\mu\text{l}$  that caused 100% mortality of the insect larvae studied after 24 hours of exposure. The extracted essential oil was classified as highly active with a lethal concentration of 46.18  $\text{mg L}^{-1}$ . Biological assays with nauplii of *Artemia salina* were performed at different concentrations of the oil to evaluate toxicity. The calculated LC50 resulted in 532.34  $\text{mg L}^{-1}$  configuring the non-toxic character, which demonstrates the safety of this essential oil for non-target organisms that coexist with the vector.

For future studies, the chemical composition of the essential oil and assays in endemic areas are necessary to identify the major metabolic species present in the botanical species, to explore its biological potential and to make viable the

development of natural larvicidal insecticides derived from the essential oil of citronella grass.

## ACKNOWLEDGMENTS

The authors express their gratitude to Pró-Reitoria de Extensão e Cultura (PROEC/UFMA) for the granting of scholarships, to Dr. V. E. Mouchrek Filho, Professor of the Department of Chemical Technology, Federal University of Maranhão, for the laboratory facilities provided, to members of the Laboratory of Research and Application of Essential Oils (LOEPAV/UFMA) for cooperation and Dr. A. Z. Silva for identifying the leaves of *Cymbopogon winterianus*.

## REFERENCES

- AGNOLIN, C. A.; OLIVO, C. J.; PARRA, C. L. C.; AGUIRRE, P. F.; BEM, C. M. D.; ZENI, D.; MOREL, A. F. Eficácia acaricida do óleo de citronela contra o *Rhipicephalus (Boophilus) microplus*. **Revista Brasileira de Saúde e Produção Animal**, v. 15, n. 3, p. 604-612, 2014.
- AMER, A.; MEHLHORN, H. Larvicidal effects of various essential oils against *Aedes*, *Anopheles*, and *Culex* larvae (Diptera, Culicidae). **Parasitology Research**, v. 99, n. 4, p. 466-472, 2006.
- BLANK, A. F.; COSTA, A. G.; ARRIGONI-BLANK, M. D. F.; CAVALCANTI, S. C.; ALVES, P. B.; INNECCO, R.; EHLERT, P. A. D.; SOUSA, I. F. D. Influence of season, harvest time and drying on Java citronella (*Cymbopogon winterianus* Jowitt) volatile oil. **Revista Brasileira de Farmacognosia**, v. 17, n. 4, p. 557-564, 2007.
- BRASIL. **Farmacopeia Brasileira**. 6. ed. Volume 1. Agência Nacional de Vigilância Sanitária. Brasília: Anvisa. 2019.
- BRASILEIRO, B. G.; PIZZILOLO, V. R.; RASLAN, D. S.; JAMAL, C. M.; SILVEIRA, D. Antimicrobial and cytotoxic activities screening of some Brazilian medicinal plants used in Governador Valadares district. **Revista Brasileira de Ciências Farmacêuticas**, v. 42, n. 2, p. 195-202, 2006.
- CANTANE, D. R.; CRISTINO, A. C.; OLIVEIRA, R. A.; FLORENTINO, H.; SANTOS, F. L.; FERNANDES, M. A.; SOUZA NETO, J. A.; RIBOLLA, P. O desenvolvimento da população do *Aedes aegypti* aplicado ao modelo de otimização no controle da Dengue. **XLVII SBPO-Simpósio Brasileiro de Pesquisa Operacional**, 2015.
- CARVALHO, A. F. U.; MELO, V. M. M.; CRAVEIRO, A. A.; MACHADO, M. I. L.; BANTIM, M. B.; RABELO, E. F. Larvicidal activity of the essential oil from *Lippia sidoides* Cham. against *Aedes aegypti* Linn. **Memórias do Instituto Oswaldo Cruz**, v. 98, n. 4, p. 569-571, 2003.

- CASSEL, E.; VARGAS, R. M. F.; MARTINEZ, N.; LORENZO, D.; DELLACASSA, E. Steam distillation modeling for essential oil extraction process. **Industrial crops and products**, v. 29, n. 1, p. 171-176, 2009.
- CAVALCANTI, E. S. B.; MORAIS, S. M. D.; LIMA, M. A. A.; SANTANA, E. W. P. Larvicidal activity of essential oils from Brazilian plants against *Aedes aegypti* L. **Memórias do Instituto Oswaldo Cruz**, v. 99, n. 5, p. 541-544, 2004.
- CHENG, S. S.; CHANG, H. T.; CHANG, S. T.; TSAI, K. H.; CHEN, W. J. Bioactivity of selected plant essential oils against the yellow fever mosquito *Aedes aegypti* larvae. **Bioresource Technology**, v. 89, n. 1, p. 99-102, 2003.
- CHENG, S. S.; LIU, J. Y.; TSAI, K. H.; CHEN, W. J.; CHANG, S. T. Chemical composition and mosquito larvicidal activity of essential oils from leaves of different *Cinnamomum osmophloeum* provenances. **Journal of Agricultural and Food Chemistry**, v. 52, n. 14, p. 4395-4400, 2004.
- CHUNG, I. M.; SEO, S. H.; KANG, E. Y.; PARK, S. D.; PARK, W. H.; MOON, H. I. Chemical composition and larvicidal effects of essential oil of *Dendropanax morbifera* against *Aedes aegypti* L. **Biochemical Systematics and Ecology**, v. 37, n. 4, p. 470-473, 2009.
- CORRÊA, J. C. R.; SALGADO, H. R. N. Atividade inseticida das plantas e aplicações: revisão. **Revista Brasileira de Plantas Mediciniais**, v. 13, n. 4, p. 500-506, 2011.
- CRUZ, R. C. D.; SILVA, S. L. C. E.; SOUZA, I. A.; GUALBERTO, S. A.; CARVALHO, K. S.; SANTOS, F. R.; CARVALHO, M. G. Toxicological evaluation of essential oil from the leaves of *Croton argyrophyllus* (Euphorbiaceae) on *Aedes aegypti* (Diptera: Culicidae) and *Mus musculus* (Rodentia: Muridae). **Journal of medical entomology**, v. 54, n. 4, p. 985-993, 2017.
- DA CRUZ, T. P.; ALVES, F. R.; MENDONÇA, R. F.; COSTA, A. V.; DE JESUS JUNIOR, W. C.; PINHEIRO, P. F.; MARINS, A. K. Atividade fungicida do óleo essencial de *Cymbopogon winterianus* jowit (Citronela) contra *Fusarium solani*. **Bioscience Journal**, v. 31, n. 1, 2015.
- DOLABELLA, M. F. **Triagem in vitro para a atividade antitumoral e anti-T. cruzi de extratos vegetais, produtos naturais e sintéticos**. 1997. Tese de Doutorado. tese de mestrado, Belo Horizonte, Universidade Federal de Minas Gerais.
- EDEN, W. T.; ALIGHIRI, D.; CAHYONO, E.; SUPARDI, K. I.; WIJAYATI, N. Fractionation of Java citronella oil and citronellal purification by batch vacuum fractional distillation. In: **IOP Conference Series: Materials Science and Engineering**. Indonesia, 2018. p. 012067.
- FALCÃO L. **Experimentais, em diferentes modelos**. Universidade Regional Integrada do Alto Uruguai e das Missões URI - Campus de Erechim. Departamento de Ciências Biológicas Programa de Pós-Graduação em Ecologia.
- FANSIRI, T.; THAVARA, U.; TAWATSIN, A.; KRASAESUB, S.; SITHIPRASASNA, R. Laboratory and semi-field evaluation of mosquito dunks® against *Aedes aegypti* and *Aedes albopictus* larvae (Diptera: Culicidae). **Southeast Asian journal of tropical medicine and public health**, v. 37, n. 1, p. 62, 2006.

FARPEJ. **Armadilha letal para mosquitos, temperada com atitude de civilidade**. 2013. Disponível em: <http://www.faperj.br/downloads/mosquiterica.pdf>. Acesso em: 30 jan. 2020.

FERREIRA, M. D. S.; BATISTA, E. K. F., DOS SANTOS FARIAS, I., Santos, L. F., de Oliveira, J. M. G., & de Sousa Silva, S. M. M. Avaliação fitoquímica e toxicológica dos extratos do fruto de *Buchenavia* sp. **Acta Brasiliensis**, v. 1, n. 2, p. 17-22, 2017.

FAIAZ, M.; MARTÍNEZ, L. C.; PLATA-RUEDA A.; GONÇALVES, W. G.; DE SOUSA, D. L. L.; COSSOLIN, J. F. S.; SERRÃO, J. E. Pyriproxyfen, a juvenile hormone analog, damages midgut cells and interferes with behaviors of *Aedes aegypti* larvae. **PeerJ**, v. 7, p. e7489, 2019.

EPA - Environmental Protection Agency. **FEDERAL INSECTICIDE, FUNGICIDE, AND RODENTICIDE ACT (FIFRA) AND FEDERAL FACILITIES**. Disponível em: <https://www.epa.gov/enforcement/federal-insecticide-fungicide-and-rodenticide-act-fifra-and-federal-facilities>. Acesso em: 10 abr. 2020.

GOMES, P. R. B.; SILVA, A. L. S.; PINHEIRO, H. A.; CARVALHO, L. L.; LIMA, H. S.; SILVA, E. F.; OLIVEIRA, M. B. Avaliação da atividade larvicida do óleo essencial do *Zingiber officinale* Roscoe (gengibre) frente ao mosquito *Aedes aegypti*. **Revista brasileira de plantas medicinais**, v. 18, n. 2, p. 597-604, 2016.

GOMES, P. R. B.; OLIVEIRA, M. B.; DE SOUSA, D. A.; DA SILVA, J. C.; FERNANDES, R. P.; LOUZEIRO, H. C.; DE PAULA, M. L.; MOUCHREK, V. E.; Fontenele, M. A. Larvicidal activity, molluscicide and toxicity of the essential oil of *Citrus limon* peels against, respectively, *Aedes aegypti*, *Biomphalaria glabrata* and *Artemia salina*. **Eclética Química Journal**, v. 44, n. 4, p. 85-95, 2019.

GOVINDARAJAN, M.; SIVAKUMAR, R.; RAJESWARI, M.; YOGALAKSHMI, K. Chemical composition and larvicidal activity of essential oil from *Mentha spicata* (Linn.) against three mosquito species. **Parasitology research**, v. 110, n. 5, p. 2023-2032, 2012.

ISMÁN, M. B. Pesticides based on plant essential oils: phytochemical and practical considerations. In: **Medicinal and Aromatic Crops: Production, Phytochemistry, and Utilization**. American Chemical Society, 2016. p. 13-26.

KAKARAPARTHI, P. S.; SRINIVAS, K. V. N. S.; KUMAR, J. K.; KUMAR, A. N.; RAJPUT, D. K.; SARMA, V. U. M. Variation in the essential oil content and composition of *Citronella* (*Cymbopogon winterianus* Jowitt.) in relation to time of harvest and weather conditions. **Industrial Crops and Products**, v. 61, p. 240-248, 2014.

LEE, HS. Mosquito larvicidal activity of aromatic medicinal plant oils against *Aedes aegypti* and *Culex pipiens pallens*. **Journal of the American Mosquito Control Association**, v. 22, n. 2, p. 292-295, 2006.

LEITE, B. L.; SOUZA, T. T.; ANTONIOLLI, A. R.; GUIMARÃES, A. G.; SIQUEIRA, R. S.; QUINTANS, J. S.; BONJARDIM, L. R.; ALVES, P. B.; BLANK, A. F.; BOTELHO, M. A.; ALMEIDA, J. R. G. S.; LIMA, J. T.; ARAÚJO, A. A. S.; QUINTANS-JUNIOR, L. J. Volatile constituents and behavioral change induced by *Cymbopogon winterianus* leaf essential oil in rodents. **African Journal of Biotechnology**, v. 10, n. 42, p. 8312-8319, 2011.

LIMA-CAMARA, T. N. Arboviroses emergentes e novos desafios para a saúde pública no Brasil. **Revista de Saúde Pública**, v. 50, p. 36, 2016.

MACEDO, I. T.; BEVILAQUA, C. M.; OLIVEIRA, L. M.; CAMURÇA-VASCONCELOS, A. L.; VIEIRA, L. D. S.; OLIVEIRA, F. R.; QUEIROZ-JÚNIOR, E. M.; TOMÉ, A. R.; NASCIMENTO, N. R. Anthelmintic effect of *Eucalyptus staigeriana* essential oil against goat gastrointestinal nematodes. **Veterinary Parasitology**, v. 173, n. 1-2, p. 93-98, 2010.

MANZOOR, F.; SAMREEN, K. B.; PARVEEN, Z. Larvicidal activity of essential oils against *Aedes aegypti* and *Culex quinquefasciatus* larvae (Diptera: Culicidae). **J. Anim. Plant Sci**, v. 23, n. 2, p. 420-424, 2013.

MENDONÇA, F. A.; SILVA, K. F. S.; SANTOS, K. K.; RIBEIRO JÚNIOR, K. A. L.; SANT'ANA, A. E. G. Activities of some Brazilian plants against larvae of the mosquito *Aedes aegypti*. **Fitoterapia**, v. 76, n. 7-8, p. 629-636, 2005.

MEYER, B. N.; FERRIGNI, N. R.; PUTNAM, J. E.; JACOBSEN, L. B.; NICHOLS, D. J.; MCLAUGHLIN, J. L. Brine shrimp: a convenient general bioassay for active plant constituents. **Planta medica**, v. 45, n. 05, p. 31-34, 1982.

MORAIS, S. M.; CAVALCANTI, E. S.; BERTINI, L. M.; BERTINI, C. L. L.; RODRIGUES, J. R. B.; CARDOSO, J. H. L. Larvicidal activity of essential oils from Brazilian *Croton* species against *Aedes aegypti* L. **Journal of the American Mosquito Control Association**, v. 22, n. 1, p. 161-164, 2006.

PEREIRA, G. P.; PRADO, A. E.; CARVALHO, R. I. N. Variação mensal do rendimento de óleo essencial de citronela. **Revista Eletrônica Científica da UERGS**, v. 2, p. 183-189, 2016.

PERTIWI, G. P. **Pengaruh Lama Waktu Penyimpanan dan Penyinaran Cahaya Terhadap Sitronelal serta Uji Toksisitas Menggunakan Metode BSLT (Brine Shrimp Lethality Test)**. 2014. Tese de Doutorado. Universitas Brawijaya.

PHASOMKUSOLSIL, S.; SOONWERA, M. Potential larvicidal and pupacidal Activities of herbal essential oils against *Culex quinque fasciatus* Say and *Anopheles minimus* (Theobald). **Southeast Asian J Trop Med Public Health**, v.41, n.6, p.1342 – 1351, 2010.

PITASAWAT, B.; CHAMPAKAEW, D.; CHOOCOTE, W.; JITPAKDI, A.; CHAITHONG, U.; KANJANAPOTHI, D.; RATTANACHANPICHAI, E.; TIPPAWANGKOSOL, P.; RIYONG, D.; TUETUN, B.; CHAIYASIT, D. Aromatic plant-derived essential oil: an alternative larvicide for mosquito control. **Fitoterapia**, v. 78, n. 3, p. 205-210, 2007.

PUSHPANATHAN, T.; JEBANESAN, A.; GOVINDARAJAN, M. Larvicidal, ovicidal and repellent activities of *Cymbopogon citratus* Stapf (Graminae) essential oil against the filarial mosquito *Culex quinquefasciatus* (Say) (Diptera: Culicidae). **Trop Biomed**, v. 23, n. 2, p.p 208-212, 2006.

ROCHA, S.; MING, L. C.; MARQUES, M. Influência de cinco temperaturas de secagem no rendimento e composição do óleo essencial de citronela (*Cymbopogon winterianus* Jowitt). **Revista Brasileira de Plantas Mediciniais**, p. 73-78, 2000.



RODRIGUES, K. A. D. F.; DIAS, C. N.; AMARAL, F. M. M.; MORAES, D. F.; MOUCHREK FILHO, V. E.; ANDRADE, E. H. A.; MAIA, J. G. S. Molluscicidal and larvicidal activities and essential oil composition of *Cymbopogon winterianus*. **Pharmaceutical biology**, v. 51, n. 10, p. 1293-1297, 2013.

SINHA, S.; JOTHIRAMAJAYAM, M.; GHOSH, M.; MUKHERJEE, A. Evaluation of toxicity of essential oils palmarosa, citronella, lemongrass and vetiver in human lymphocytes. **Food and Chemical Toxicology**, v. 68, p. 71-77, 2014.

TABANCA, N.; ÖZEK, G.; ALI, A.; DURAN, A.; HAMZAOGLU, E.; BASER, K. H. C.; KHAN, I. A. Chemical composition of *Heracleum pastinacifolium* subsp. *transcaucasicum* and subsp. *incanum* essential oils, and their biting deterrent and larvicidal activity against *Aedes aegypti*. **Planta Medica**, v. 78, n. 05, p. P\_89, 2012.

TAWATSIN, A.; WRATTEN, S. D.; SCOTT, R. R.; THAVARA, U.; TECHADAMRONGSIN, Y. Repellency of volatile oils from plants against three mosquito vectors. **Journal of vector ecology**, v. 26, p. 76-82, 2001.

VELETOVAC, A. **Toxic essential oils**. Tese (Doutorado em Farmácia) – Universität Wien. Viena, p 59. 2016.

VELOSO, R. A.; CASTRO, H. G.; CARDOSO, D. P.; CHAGAS, L. F. B.; FREITAS, A.; JÚNIOR, C. Óleos essenciais de manjeriço e capim citronela no controle de larvas de *Aedes aegypti*. **Revista Verde (Pombal-PB-Brasil)** v. 10, n. 2, p. 101-105, 2015.

WHO. **World Health Organization Technical Reports Series n. 443**, 1970.

WOLFFENBUTTEL, A. N. Óleos essenciais. **Informativo CRQ-V**, ano XI, n. 105, págs. v. 6, 2007.

ZARA, A. L. D. S. A.; SANTOS, S. M. D.; FERNANDES-OLIVEIRA, E. S.; CARVALHO, R. G.; COELHO, G. E. Estratégias de controle do *Aedes aegypti*: uma revisão. **Epidemiologia e Serviços de Saúde**, v. 25, p. 391-404, 2016.